



**Fig. 1.** (A) Homocysteine and (B) vascular endothelial growth factor 2 (VEGFR-2) serum levels in control, hemodialyzed (HD), and kidney transplanted (KTx) patients. \* $P < 0.05$  vs. KTx patients; \*\* $P < 0.001$  vs. KTx and control patients; # $P < 0.001$  vs. KTx patients and  $P < 0.014$  vs. control patients.

the renin-angiotensin system. Therefore, specific dosage of ACE-I and ARBs may be required to obtain maximal renoprotective effects in various populations. Unfortunately, the optimal dosage of ACE-I has only been investigated in Japanese patients with nonrenal disease [1].

PETER JACOBSEN and HANS-HENRIK PARVING  
Gentofte, Denmark

Correspondence to Peter Jacobsen, Steno Diabetes Center, Niels Steensens Vej 2, 2820 Gentofte, Denmark.  
E-mail: pkjacobsen@dadlnet.dk

## REFERENCES

1. NAKAO N, YOSHIMURA A, MORITA H, *et al*: Combination treatment of angiotensin-II receptor blocker and angiotensin-converting-enzyme inhibitor in non-diabetic renal disease (COOPERATE): A randomised controlled trial. *Lancet* 361:117–124, 2003
2. JACOBSEN P, ANDERSEN S, ROSSING K, *et al*: Dual blockade of the renin-angiotensin system versus maximal recommended dose of ACE inhibition in diabetic nephropathy. *Kidney Int* 63:1874–1880, 2003
3. EXNER DV, DRIES DL, DOMANSKI MJ, COHN JN: Lesser response to angiotensin-converting-enzyme inhibitor therapy in black as compared with white patients with left ventricular dysfunction. *N Engl J Med* 344:1351–1357, 2001

## Inverse association between homocysteine and vascular endothelial growth factor receptor 2 serum levels in hemodialyzed and kidney transplanted patients

**To the Editor:** Homocysteine has an inhibitory effect on angiogenesis [1]. However, the vascular endothelial

growth factor receptor-2 (VEGFR-2) gene has recently been found up-regulated on human atherosclerotic plaques [2]. In order to analyze the association between homocysteine and VEGFR-2 serum levels, we have randomly selected 26 regular hemodialysis (HD) patients ages  $56 \pm 12$  years (mean  $\pm$  SD; M = 14, F = 12), and 26 age-matched kidney transplanted individuals (KTx), ages  $52 \pm 8$  years (M = 21, F = 5). KTx transplant age was  $13 \pm 9$  months and creatinine was  $1.6 \pm 0.5$  mg/dL. Nine age-matched healthy volunteers ( $47 \pm 5$  years) were also recruited. HD samples were collected at the end of the long interval period. Serum homocysteine (Hcy) and VEGFR-2 were tested by high-performance liquid chromatography (HPLC) with fluorimetric detection and a commercial enzyme-linked immunosorbent assay (ELISA) kit, respectively. Hcy was higher in HD patients ( $57 \pm 36$   $\mu\text{mol/L}$ ) than in KTx patients ( $23 \pm 8$   $\mu\text{mol/L}$ ;  $P < 0.001$ ) and control patients ( $15 \pm 6$   $\mu\text{mol/L}$ ;  $P < 0.01$  vs. HD;  $P < 0.05$  vs. KTx). VEGFR-2 was lower in HD patients ( $1705 \pm 387$   $\text{pg/mL}$ ) than in KTx patients ( $2449 \pm 825$   $\text{pg/mL}$ ;  $P = 0.014$ ) or control patients ( $2150 \pm 464$   $\text{pg/mL}$ ) (Fig. 1). The present study shows that Hcy and circulating VEGFR-2 are inversely associated, consistent with previous reports showing that homocysteine inhibits angiogenesis. Our data also suggest that up-regulation of the VEGFR-2 gene is probably a local finding, and that hyperhomocysteinemia may be involved in the release of VEGFR-2 into the systemic circulation.

VINCENZO SEPE, CARMELO LIBETTA, NATALIA ROSSI,  
CRISTINA GUIDETTI, and ANTONIO DAL CANTON  
Pavia, Italy

Correspondence to Vincenzo Sepe, Unit of Nephrology, Dialysis and Transplantation, IRCCS Policlinico San Matteo, University of Pavia, Italy.  
E-mail: vsepe@libero.it

## REFERENCES

1. HAJJAR KA: Homocysteine: A sulph'rous fire. *J Clin Invest* 107:663–664, 2001
2. HILTUNEN MO, TUOMISTO TT, NIEMI M, *et al*: Changes in gene expression in atherosclerotic plaques analyzed using DNA array. *Atherosclerosis* 165:23–32, 2002